



Biological protection against grape berry moths. A review

Denis Thiéry^{1,2} · Philippe Louâpre³ · Lucile Muneret¹ · Adrien Rusch¹ · Gilles Sentenac⁴ · Fanny Vogelweith⁵ · Corentin Iltis³ · Jérôme Moreau³

Accepted: 16 February 2018

© INRA and Springer-Verlag France SAS, part of Springer Nature 2018

Abstract

Grape is a major crop, covering 7.5 M ha worldwide, that is currently being confronted with three main challenges: intensive pesticide use that must be reduced, invasion by new pests/diseases, and climate change. The biological control of pests and vectors would help address these challenges. Here, we review the scientific literature on the biological control of grape moths by macroorganisms (excluding nematodes). Two components, biological control with an active human role, mainly using biocontrol agents through inundation or inoculation, and conservation biological control, are considered. The major points are the following. (1) Tortricid grape moths seriously damage grapes worldwide, causing yield losses and quality reduction. The more geographically widespread species, *Lobesia botrana*, continues to extend its range, invading South American and, more recently, North American vineyards. (2) Parasitoids and predators (including arthropods, birds, and bats) that can control grape pests are very diverse. (3) Different methods exist to assess pest control efficiency in the field but some of them remain to be developed. (4) Environmental factors, including host plants, landscape, grass or floral covers, and organic practices, affect the natural control of grape moths. (5) Pest resistance to parasitoids strongly depends on their immune system, which is controlled by the host plant. Future climate changes may modify this tritrophic interaction and thus affect biological control strategies. We conclude that biological control has a great deal of potential in viticulture and that addressing these key factors would improve the efficiency levels of biological control strategies. This would help growers and stakeholders to significantly reduce insecticide use in vineyards.

Keywords *Lobesia botrana* · *Eupoecilia ambiguella* · Biodiversity · Agroecology · Viticulture · Agricultural practices · Landscape architecture · Parasitoids · Predators

Contents

1. Introduction
 2. Grape berry moths in vineyards
 3. Grape pests' natural enemies in vineyards
 - 3.1 Parasitoids
 - 3.2 Arthropod predators
 - 3.3 Predation by birds and bats
 4. Evaluation of pest control efficiency in the field
 5. Key factors inducing variation in biological control of grapevine moths
 - 5.1 Effect of the host plant (including grape cultivar)
 - 5.2 Viticulture practices and biological control
 - 5.3 Vineyard landscape and natural biological control
 - 5.4 Climate changes on tritrophic interactions and natural biological control
 6. Perspectives and conclusions
 - 6.1 Challenges for research
 - 6.2 Challenges for growers and crop protection advisors
- Acknowledgements
References

✉ Denis Thiéry
denis.thiery@inra.fr

¹ INRA UMR 1065 Santé et Agroécologie du Vignoble, INRA, ISVV, Villenave d'Omon Cedex, France

² UMT Seven INRA-IFV, Villenave d'Omon Cedex, France

³ Equipe Ecologie Evolutive, UMR CNRS 6282 Biogéosciences, Université Bourgogne Franche-Comté, Dijon, France

⁴ IFV Beaune, 6, rue du 16è Chasseurs, 21200 Beaune, France

⁵ Institut für Zoologie, Abt. Evolutionsbiologie, Johannes Gutenberg-Universität Mainz, Johannes-von-Müller-Weg 6, 55128 Mainz, Germany

1 Introduction

Crops worldwide are the targets of more than 10,000 insect species (Dhaliwal et al. 2010), which are responsible for huge annual losses of between 20 and 50% of total production (Thacker 2002; Oerke 2006). Given the rate of human population growth, protecting crops from damage caused by pests is a major challenge to ensuring an adequate food supply in the future (Thomas 1999; Tilman et al. 2011). The extensive use of chemical products to control crop pests is now widely criticized because of their negative effects on the environment and trophic chains, from plants to humans (Hallenbeck and Cunningham-Burns 1995; Desneux et al. 2007; Geiger et al. 2010). Moreover, owing to the past irrational use of pesticides, some insect pests have acquired forms of resistance, making them less susceptible to biochemical products (Roush and Tabashnik 1990; Boyer et al. 2012). For the past few decades, the scientific community and governmental agencies have recommended the use of alternative, less invasive and more environmentally friendly methods of control to make agriculture more sustainable. Controlling pests through biological control and integrated pest management strategies are potential ways to address current ecological and societal concerns (Kogan 1998; Thomas 1999; Way and Van Emden 2000; Brewer and Goodell 2012).

Biological control and integrated pest management aim at optimally managing insect pests using different techniques, like the use of chemical mediators to disrupt mating (i.e., insect pheromones), autocidal control (based on the release of sterile males), and auxiliary macroorganisms (Thacker 2002; Walter 2005), without disturbing ecosystem functions (Ehler 2006; Naranjo and Ellsworth 2009; Abrol and Shankar 2012). At least two strategies based on the activity of auxiliary macroorganisms can be developed: (i) biological control based on endogenous biodiversity from crops and the adjacent seminatural habitat and (ii) natural enemy releases. To maximize their efficiency, such control methods must consider the ecology of targeted pests and natural enemies, as well as the relationships among trophic levels, including host plants (Ferguson et al. 2005; Pérez-Staples et al. 2012).

Here, we focus on grape, which is a major crop worldwide, with a vine area of 7.52 million ha that is targeted by numerous pests and diseases, resulting in very high levels of pesticide treatments (e.g., in 2010, the treatment frequency ranged from 10 to 25 times per year in France; Butault et al. 2010) with total amounts of active substance in kilograms per hectare in average 21.4 (EU) and, respectively, 49.6 (Portugal), 32.6 (France), 31.1 (Germany), 20.3 (Greece), and 17.8 (Italy) (Endure 2006). Among the numerous insect pests of grapes, the two tortricid Lepidoptera, the European grapevine moth *Lobesia botrana* (Den. & Schiff.) and the grape berry moth (GBM) *Eupoecilia ambiguella* (Hubn.), have been worldwide harmful pests for years in viticulture and often

cause high levels of damage to grape bunches (Thiéry et al. 2011a) (Fig. 1). These two main species have very similar traits and will be grouped further in this review as GBMs. Surprisingly, no accurate quantitative economic impact is available for GBMs. One *L. botrana* larva is capable of damaging between 2 and 10 berries, depending on the cultivar and the grape phenology, and up to 20 to 30 larvae per cluster may occur in heavily attacked vineyards (Delbac and Thiéry 2016). In addition to direct predation on the berry, the presence of larvae encourages bunch rot development (causal agents being *Botrytis cinerea*, *Aspergillus carbonarius*, and *Aspergillus niger*), which results in severe qualitative and quantitative damages (Cozzi et al. 2006; Delbac and Thiéry 2016). Currently, these pests are mainly controlled by synthetic insecticides, which can be neurotoxic (e.g., pyrethrinoids and indoxacarb); insect growth regulators, like flufenoxuron; and recently, muscle contraction inhibitors, like emamectin. Thus, viticulture must be adapted to the new challenges of pest management, and the development of biological controls against these pests is promising (Thiéry 2011). Alternative treatments against GBMs have already been produced, and they mainly rely on natural insecticides including sprays of *Bacillus thuringiensis* (Bt) toxin (see Pertot et al. 2017 for a recent review) and on mating disruption by sex pheromones (recently reviewed in Ioriatti et al. 2011). Biological control based on macroorganisms (parasitoids and predators) could also be developed as a valuable alternative to chemical pest control in viticulture in combination with these two methods (Mills and Danne 2005; Moreau et al. 2009; Thiéry 2011).




Fig. 1 Fully grown (5th instar) larva of **a** *Lobesia botrana* (picture F. Vogelweith) and **b** *Eupoecilia ambiguella* (picture F. Vogelweith), on vine leaves. Larva length is c. 1 cm

Despite the numerous studies carried out in ecological chemistry, physiology, and behavior to improve the effectiveness of grape pest management (Thiéry 2008; Ioriatti et al. 2011), GBMs continue to proliferate in vineyards, which suggests that control methods have not been optimized.

The objective of this review is to synthesize the literature regarding the biological control of GBMs. It will focus on the biological control of GBMs by beneficial macroorganisms, including arthropods, birds, and bats, and will consider the biological and ecological factors that may affect the efficiency levels of such biological control strategies. The key factors identified in this review should help increase our understanding of the context dependency and the variability associated with biological control strategies and thus aid in their future development to protect grapes from GBMs.

2 Grape berry moths in vineyards

The two GBMs, *L. botrana* and *E. ambiguella*, are polyphagous and plurivoltine, producing two to four generations per year under European vineyard conditions (Thiéry 2008) (Fig. 2). While the larvae develop on more than 30 different food plants, these polyphagous species occur mainly on grape varieties and almost exclusively develop on flowers and berries. Therefore, grape provides an important and continuous resource from spring to autumn, allowing the establishment of high population levels, which can reach over 15–30 larvae per

grape cluster in some places (Thiéry et al. 2014; Schellhorn et al. 2015; Delbac and Thiéry 2016). They undergo diapause in the winter as pupae under grape stock bark or inside the nonharvested grape bunches, and then adults emerge in early spring for nuptial flights. Interestingly, *L. botrana* and *E. ambiguella* belong to the Olethreutini subfamily, a group of species that oviposit individual eggs unlike most moths, which lay egg clusters (Moreau et al. 2016). This trait is interpreted as an oviposition strategy for limiting exposure to egg parasitoids and predators, which have a discrete and cryptic host/prey resource. Neonatal larvae can penetrate berries within the few hours after hatching. Fully grown larvae rarely move from one bunch to another (Torres-Vila et al. 1997), and when the adult density is high, high larval populations may also occur, with up to 30 larval *L. botrana* per *Cabernet sauvignon* bunch (Thiéry et al. 2014). Thus, the spatial distributions of eggs and larvae are mostly the result of the oviposition behavior performed by the females (Gabel and Thiéry Gabel and Thiéry 1992, 1996). During their development in June, the larvae of these two moth species build individual nests, called “glomerulae,” with their silk (Delbac and Thiéry 2016). The nests provide shelter and protection against adverse conditions (i.e., insulation from temperature variation and a barrier against predators and parasitoids). In July and August, the larvae of these two species are berry borers.

3 Grape pests' natural enemies in vineyards

Globally, both the species richness and evenness of natural enemies increase the capabilities of natural pest control services in agrosystems (Letourneau et al. 2009; Crowder et al. 2010). This is the result of nonexcluding mechanisms, either a complementarity effect or a sampling effect (Straub et al. 2008). The complementarity effect predicts that pest suppression resulting from different natural enemy species is equal or greater than the sum of the suppression induced by each species alone. This relationship can be explained by resource complementarity owing to niche partitioning or facilitation (Straub et al. 2008). The sampling effect hypothesis states that a larger number of species in a given assemblage increase the probability of including an effective predator that contributes more than the other to the pest control function (Loreau and Hector 2001; Straub et al. 2008). Promoting the abundance and diversity of natural enemies is therefore expected to optimize natural pest control services in vineyard landscapes. However, the exact roles of several species and functional groups found in vineyards are not known.

Contributions to the biological control of GBM can come from either natural populations of arthropod predators or parasitoids and also from avian and mammalian predators (Thiéry et al. 2001; Sentenac 2011; Vincent et al. 2012; Rusch et al.




Fig. 2 Life cycle of grape berry moths, as represented for *L. botrana* showing the main stages. Duration is indicative and varies as a function of external temperature. Durations provided are those recorded in vineyards for the spring generation in south west France (1 month in summer). The two species *L. botrana* and *E. ambiguella* have very similar life cycle. In SW France, *Lb* accomplishes three or four generations per year, *Ea* two or three cycles (D. Thiéry and N. Maher)

2015 for a recent issue). Increasing attention is currently being placed on arthropod biodiversity, which may have potential owing to their natural predation and parasitism capabilities. As examples, impressive numbers of carabid beetle species (124 in Goulet et al. 2004; 39 in Rusch et al. 2016) and 97 spider species (Bolduc et al. 2005) have been found in vineyards. Larval or egg parasitoids are also very abundant and diverse in vineyards (Thiéry et al. 2006; Thiéry et al. 2011b; Moreau et al. 2009; Sentenac 2011). To date, these studies have only established the occurrence of a large diversity of natural enemies, but their control potential has not been evaluated.

3.1 Parasitoids

Current vineyards are far from being “no parasitoid lands.” In lectures or training courses, students and growers are always surprised by a list of parasitoids found in most vineyards, with more than 70 species that are mainly found in Hymenoptera (Thiéry 2008; Loni et al. 2016) (see Table 1 for an abbreviated list). Recently, in a study conducted in Australian vineyards, 20–30 parasitoid species were found (Paull and Austin 2007). In addition to the host density, which is a main driver of parasitoid populations, viticulture practices, such as organic viticulture or the use of interrow grass covers and floral strips, also favor a complex parasitoid community (Genini 2000). Globally, recent research indicates that landscape and climatic conditions also affect the diversity of parasitoids that naturally occur in a vineyard (Moreau et al. 2009; Loni et al. 2016). Clearly, understanding how such factors contribute to the variation in biodiversity requires further study.

Some species of parasitoids are rather cosmopolitan and found worldwide, such as the Ichneumonid *Campoplex capitator* or the Pteromalids *Dibrachys* spp., possibly because they are more plastic in their ecological requirements (Moreau et al. 2010). Other species are less frequently observed and are limited to certain vineyards or vine production regions. An example is *Exochus tibialis*, which was exclusively found in Switzerland (in the Valaisan vineyards, Switzerland) and Alsace (France) in a large field survey performed by Moreau et al. (2009).

The Ichneumonid *C. capitator* (previously named *Campoplex majalis* for its strong occurrence in spring) was probably the first well-studied larval parasitoid (Audouin 1842; Jolicoeur 1894). This species is described as a specialist parasitoid that has a large expected foraging distance (Xuéreb and Thiéry 2006). Because it is able to undergo diapause in its host, it has the advantage of good synchrony with the pest and is thus very active in the first spring generation.

In addition to Hymenoptera, the tachinids from the order Diptera can be important parasitoids, and *Phytomyia nigrina* is suspected to increase its geographical area in the context of climate change (Reineke and Thiéry 2016). *P. nigrina* is an efficient parasitoid in warm vineyards

(Thiéry et al. 2006) and will represent an interesting larval parasitoid against *L. botrana* in future years.

Trichogramma sp. has been known egg parasitoids in viticulture for more than a century. For instance, *Oophthora semblidis* (current name *Trichogramma semblidis*) was found in *L. botrana* and *E. ambiguella* eggs from several vineyards (Marchal and Feytaud 1911). To date, only the release of *Trichogramma* sp. has been attempted in viticulture on a large scale. The main reasons being that the moth eggs are easily mass produced and inexpensive, and the resulting larvae provide an efficient level of biocontrol against other moth pests (e.g., the European corn borer). They were first used in 1980 against GBM eggs (Barnay et al. 1999; Reda Abd el Monsef 2004; Hommay et al. 2011; Walton et al. 2012), the American grape berry moth *Endopiza viteana* (Nagarkatti et al. 2003), and also against the light brown apple moth *Epiphyas postvittana* in Australian vineyards (Glenn and Hoffmann 1997). Several species of *Trichogramma*, such as *T. brassicae* Bezdenko, *T. cacoeciae* Marchal, *T. dendrolimi* Matsumura, and *T. minutum* Riley, have been used with significant but varying pest reduction results. Currently, the use of *Trichogramma* sp. against GBMs is still very marginal in viticulture, mainly because of the varying and inconsistent efficiency levels (Barnay et al. 1999; Walton et al. 2012). Even though the use of sulfur applications is suspected as deleterious for *Trichogramma* sp., we did not identify a scientific report stating that specific conclusion. Thus, the main factors for explaining these varying parasitism rates are the amounts of *Trichogramma* sp. released per surface area and the number of release points, which affects the size of the protected area (Hommay et al. 2011). Hommay et al. (2002) reported better results against the GBM when releasing *Trichogramma evanescens* at 800 instead of 400 points per hectare.

To increase the efficiency of *Trichogramma* sp. releases, several experiments investigated the roles of kairomones in the attraction of *Trichogramma* sp. The egg odors of *L. botrana* and *E. ambiguella* have been analyzed and rely mainly on C16–C18 fatty acids and esters (Gabel and Thiéry 1992; Thiéry et al. 1995). They are well detected by conspecific females, and the arrestment of *T. brassicae* has been described in response to oleic acid (Frenoy et al. 1992). Some evidence also exists that host females (pheromone and wing scales) may attract *Trichogramma* sp. (Fatouros et al. 2008; Milonas et al. 2009). Thus, the marking of the alternative host *Ephesia kunhiella* eggs with *L. botrana* scales in the laboratory increased the oviposition rates by *T. cacoeciae* (Barnay et al. 1999). This suggests that several *Trichogramma* sp. can use such kairomones to optimize their foraging behaviors in vineyards and that such semiochemicals may be useful for improving the efficiency of biological control programs. Surprisingly, such research efforts on *Trichogramma* sp. against grape moths have not been sustained in vineyards for the last few years. The three following reasons could help in

Table 1 Nonexhaustive list of parasitoids (alphabetic order) reported from the literature as natural enemies of grape moths in vineyards in west European countries. This list is selected and implemented from Thiéry (2008) and Sentenac (2011). Only species reported by at least 2 references in the former list are presented here

Species	Family	Host orders	Reported hosts in vineyards	Parasitized instars (when known)
<i>Agrothereutes abbreviatus</i> (F.)	Ichn	Lepidoptera	EA, LB	Pupae
<i>Ascogaster quadridenata</i> (Wesm.)	Ichn	Tortricidae	LB	Larvae, pupae
<i>Brachymeria minuta</i> (Wesm.)	Chal	Lep, Dip	EA	
<i>Campoplex capitator</i> (Aubr.)	Ichn	Tortricidae	EA, LB	L3-L4
<i>Diadegma fenestrata</i> (Holm.)	Ichn	Lepidoptera	LB, SP	Larvae
<i>Dibrachys affinis</i> (Masi)	Chal	Lepidoptera and other insects	EA, LB, SP	L4-L5
<i>Dibrachys cavus</i> (Walk.) (syn <i>boucheanus</i>)	Chal	Lepidoptera and other insects	EA, LB, SP	L4-L5, pupae
<i>Dicaelotus inflexus</i> (Thom.)	Ichn	Lepidoptera	LB	Pupae
<i>Dicaelotus resplendens</i> (Holm.)	Ichn	Lepidoptera		
<i>Elachertus affinis</i> (Masi)	Chal	Tortricidae	EA, LB, SP	Larvae
<i>Exochus tibialis</i> (Holm.)	Ichn	Lepidoptera	LB	Larvae, pupae
<i>Gelis areator</i> (Panz.)	Ichn	Lep, Hym	EA, LB	Larvae
<i>Goniozus claripennis</i> (Först.)	Beth	Lepidoptera	PS	Larvae
<i>Ischnus alternator</i> (Grav.)	Ichn	Lepidoptera	LB	Pupae
<i>Itoplectis alternans</i> (Grav.)	Ichn	Lep, Hym	LB, SP	Pupae
<i>Itoplectis maculata</i> (Fabr.)	Ichn	Lep, Hym	EA, LB, SP	Pupae
<i>Itoplectis tunetana</i> (Schmied.)	Ichn	Lep, Hym	EA, LB	Pupae
<i>Phaeogenes melanogonos</i> (Gmel.)	Ichn	Lepidoptera	EA, SP	Pupae
<i>Phaeogenes planifrons</i> (Wesm.)	Ichn	Lepidoptera	SP	Pupae
<i>Phytomyza nigrina</i> (Meig.) (= <i>nitidiventris</i>)	Tach	Lepidoptera	LB	Larvae
<i>Pimpla spuria</i> (Grav.)	Ichn	Lepidoptera	LB	Pupae
<i>Pimpla turionellae</i> (L.)	Ichn	Lepidoptera	EA, LB, SP	Pupae
<i>Pteromalus</i> spp. (>8 species)	Ichn	Lep, Dip, Col, Hym	EA, LB, SP	Larvae, pupae
<i>Scambus elegans</i> (Woldst.)	Ichn	Lep, Hym	LB	Larvae
<i>Tranosemella praerogator</i> (L.)	Ichn	Lepidoptera	EA, LB, SP	Larvae
<i>Trichogramma Minutum</i> (Riley)	Chal	Lepidoptera	EA, LB, SP	Eggs
<i>Trichogramma cacoeciae</i> (Marchal)	Chal	Lepidoptera	EA, LB, SP	Eggs
<i>Trichogramma evanescens</i> (West.)	Chal	Lepidoptera	EA, LB, SP	Eggs
<i>Triclistus</i> sp.	Chal	Lepidoptera	LB	Larvae, pupae

EA, *Eupoecilia ambiguella*; LB, *Lobesia botrana*; SP, *Sparganothis pilleriana*; Beth, Hym. Bethyridae; Chal, Hym. Chalcidoidea; Ichn, Hym. Ichneumonidae; Tach, Tachinidae. *Pteromalus* and *Trichogramma* spp. regroup several species

explaining this occurrence: (i) The recent development of insect growth regulators and muscle inhibitors, which are easy to use, efficient, and cheap, in viticulture probably reduced the competitiveness of parasitoid release methods; (ii) to a minor extent, mating disruption with sexual pheromones against *L. botrana* and *E. ambiguella* also negatively affected the development of such inundation or inoculation biological control techniques; and (iii) for larval parasitoids, to our knowledge, no commercial product is available for use in vineyards. *C. capitator* has been reared (Xuéreb and Thiéry 2006), as were the two species of *Dibrachys* (*cavus* and *affinis*) (Chuche et al. 2006) but only for research purposes. We hope that the commercial production of such biocontrol agents will become feasible relatively soon.

3.2 Arthropod predators

Owing to the variable habitats occupied by tortricid moths over their life span (eggs and larvae in vegetation, flying adults and nymphs wintering under the bark or in the soil), they are exposed to numerous vertebrate and invertebrate predator species. The arthropod predators of GBMs have been classified into occasional and regular predators; however, knowledge regarding their biology and impacts on pest populations comes from other crops (Sentenac 2011). Arthropod predators that are involved in the top-down control of grape moths encompass a large range of species, including spiders (each vineyard in the Bordeaux region can shelter several dozens of species; Muneret, PhD thesis unpublished data), harvestmen, true bugs

(Miridae, Anthocoridae, Nabidae, and Reduviidae), lacewings (*Chrysopa perla*, *Chrysoperla carnea*, *Chrysoperla lucasina*, *Chrysoperla affinis*, *Dichochrysa flavifrons*, and *Dichochrysa prasina* have been found in vineyards in France), and syrphids (only one species, *Xanthandrus comtus*) (Sentenac et al. 2011). To our knowledge, the identities of the more effective predators of tortricids in vineyards have been poorly investigated, and for the majority of these groups, limited data exists on their distributions in vineyard landscapes. For instance, harvestmen have been detected as tortricid larvae consumers in French vineyards (unpublished data). Recent studies on the codling moth in apple orchards showed that ground spiders are mainly involved in the predation of emergent nymphs during spring, while carabid beetles are involved in the predation of pupae during autumn (Boreau de Roince et al. 2012). In addition, heteroptera, ants, and earwigs are involved in egg predation (Glen 1977; Glen and Milsom 1978; Frank et al. 2007). Additionally, 15% of earwigs, 8% of carabids, and 8% of spiders have ingested the tortricid codling moths (Unruh et al. 2016). Such data can be reasonably extrapolated to the vineyard system, which is similar to the orchard system. Spiders are rather diverse in vineyards (see Table 2 for the species detected in Bordeaux vineyards) and may be considered the key predators for controlling grape moths (Hogg and Daane 2010).

3.3 Predation by birds and bats

The predation of insect pests by birds and bats has been observed in several case studies (Bael et al. 2008; Karp et al. 2013; Maas et al. 2013). However, using birds and bats for biocontrol is currently being debated. Several observations indicate that birds may have adverse effects because they may also consume the grapes (Skopura and Hothem 1985; Watkins et al. 2000) and also arthropod's natural enemy (Jedlicka et al. 2014). Bats, however, would probably be very efficient, but we miss published reports in grapes and how to actively manage and use bats is less clear, especially since their spatial foraging range is not clearly known. For instance, the combined exclusion of birds and bats around cacao trees caused a significant increase in phytophagous insects and a decrease of 31% in crop yield (Maas et al. 2013). Similarly, in a coffee system, birds reduced phytophagous infestations by approximately 50% (Karp et al. 2013). These results highlight the key roles of birds and bats in the biological control of pests in perennial systems. Avian predation of the grape moth is currently motivating important research efforts (see for example Barbaro et al. 2017). The bird predation of moth larvae was affected by the interaction of bird functional community structure with habitat heterogeneity. However, to date, limited scientific literature is available on this topic. Moreover, Jedlicka et al. (2011) investigated the effects of conservation measures, such as providing nest sites for birds in vineyards. This measure strengthened pest control services

Table 2 Main arachnid species collected in pitfall traps from the east Bordeaux vineyard (Entre Deux Mers and Libourne, Bordeaux vineyards). Family ranked by alphabetic order. Data source: Lucile Muneret PhD unpublished data. Identification by morphological criteria (Roberts 2009 and <https://arachno.piwigo.com/>)

Families	Species
Agelenidae	<i>Eratigena agrestis</i>
Gnaphosidae	<i>Callilepis nocturna</i>
Gnaphosidae	<i>Civizelotes civicus</i>
Gnaphosidae	<i>Drassodes lapidosus</i>
Gnaphosidae	<i>Drassyllus praeficus</i>
Gnaphosidae	<i>Gnaphosa lucifuga</i>
Gnaphosidae	<i>Haplodrassus dalmatensis</i>
Gnaphosidae	<i>Micaria coarctata</i>
Gnaphosidae	<i>Micaria pulicaria</i>
Gnaphosidae	<i>Setaphis carmeli</i>
Gnaphosidae	<i>Trachyzelotes fuscipes</i>
Gnaphosidae	<i>Zelotes aeneus</i>
Linyphiidae	<i>Agyneta rurestris</i>
Linyphiidae	<i>Diplostyla concolor</i>
Linyphiidae	<i>Erigone dentipalpis</i>
Linyphiidae	<i>Mermessus trilobatus</i>
Linyphiidae	<i>Oedothorax apicatus</i>
Linyphiidae	<i>Tenuiphantes tenuis</i>
Liocranidae	<i>Agraecina lineata</i>
Lycosidae	<i>Arctosa perita</i>
Lycosidae	<i>Hogna radiata</i>
Lycosidae	<i>Pardosa agrestis</i>
Lycosidae	<i>Pardosa hortensis</i>
Lycosidae	<i>Pardosa prativaga</i>
Lycosidae	<i>Pardosa proxima</i>
Lycosidae	<i>Pardosa vittata</i>
Lycosidae	<i>Trochosa robusta</i>
Lycosidae	<i>Xerolycosa miniata</i>
Salticidae	<i>Neaetha membraosa</i>
Salticidae	<i>Salticus scenicus</i>
Tetragnathidae	<i>Pachygnatha degeeri</i>
Theridiidae	<i>Asagena phalerata</i>
Thomisidae	<i>Ozyptila sanctuaria</i>
Thomisidae	<i>Xysticus erraticus</i>
Thomisidae	<i>Xysticus kochi</i>
Zodariidae	<i>Zodarion italicum</i>

to vineyard growers. The presence of nest boxes increased the species richness of avian insectivorous by 50% as well as the predation rates of *L. botrana* by 2.4 times compared with controls. These results in vineyards are in agreement with the literature on the potential positive roles of birds in pest control. However, birds can also have negative effects on overall pest control services through the predation of other natural enemies (e.g., spiders and parasitoids) (Martin et al.

2013; Jedlicka et al. 2014). The specific roles of birds in the control of specific pest species, such as grape moths, in vineyards are still debated and further applied research on management options to optimize avian predation services is needed.

The predation of flying moths by bats is already known and is currently gaining research interest in vineyards (Rydell et al. 1996). Bats are nocturnal, like adult grape moths, and observations of bats flying at a 5–10-m range above the vines match the flying behavior of *L. botrana* (our unpublished data). However, very few publications have focused on the predation activity of bats in vineyards. Predation, or at least contact with prey, can be recorded by typical ultrasonic vibrations, and recent observations in Burgundy (Sentenac, unpublished data) confirm that the predation of tortricid moth by bats could be considered as a potential factor in biological control. This ongoing study has presently recorded more than 15 species of bats in Burgundy vineyards (Table 3), with the most active species being *Pipistrellus pipistrellus*, *Pipistrellus kuhlii*, and *Eptesicus serotinus*.

Predation should, however, be checked carefully. First is by confirming the amount of grape moth adults of each species predated by analyzing DNA traces in bat guano (see also Section 4). Additionally, the feeding range of bats has to be carefully studied to allow their nesting behavior to be adapted to the control of grape moths. Interestingly, dispensers of *Thaumetopoea pityocampa* pheromones placed in pine forests positively influenced bat activity (Charbonnier et al. 2014). Even though the sonic detection by GBM has not been described to date, the capacity to escape bats should, however, be considered (Speakman and Rydell 2000). The escaping flight strategies of grape moths could be a component of the zigzagging or circular flight patterns observed in males and females, respectively. This type of flight has been observed in *L. botrana* (Gabel and Thiéry 1994) but may not be related to a bat escape behavior.

4 Evaluation of pest control efficiency in the field

As detailed above, vineyards are characterized by a high, sometimes unexpected, biodiversity that contains a large number of grape moth enemies. To estimate the potential natural pest control services in a vineyard, the effective predators and parasitoids that efficiently control GBM should be identified, and the predation and parasitism rates should be quantified in the field (see Birkhofer et al. 2017 for a complete review of methods to quantify biological pest control in the field). However, assessing the level of pest control in an agroecosystem is challenging because it requires a combination of different techniques for an accurate measurement (Birkhofer et al. 2017).

The most employed methods to assess parasitism rate in the field, and thus the potential control of pests by the parasitoid community, are based on a huge sampling of hosts as larvae (e.g., Moreau et al. 2009) or the use of sentinel methods (for example, the exposure of eggs to parasitoids) (Rusch et al. 2017a). The identification of parasitoid species is mainly based on morphological traits and requires accurate skills in systematics (MacFadyen et al. 2009; Sentenac 2011; Rusch et al. 2015). Such techniques are, however, time consuming and necessitate the good handling of collected pest larvae, which have to be raised until emergence (Agusti et al. 2005; Traugott et al. 2006). It also often leads to the underestimation of biological control potential because of sample death, and these techniques are often not compatible with the rapid detection of the parasitoid (Jourdie et al. 2008; Hrcek et al. 2011; Papura et al. 2016). To counteract these problems, more recent molecular methods that enable the detection of prey-specific DNA in the host body (for parasitoids), including grape berry moths, have been developed (Birkhofer et al. 2017). Two main complementary approaches are currently used in this very fast-moving field: diagnostic PCR that uses species-specific primers and DNA barcoding that uses next-generation sequencing and universal primers. Such methods allow the assessment of parasitism rates of several species simultaneously inside collected moth larvae (Wirta et al. 2014; Papura et al. 2016). Such techniques should, however, be improved by making them simpler and cheaper so that they can be used by technicians and vineyard advisors.

Additionally, using artificial prey, such as dummy caterpillars made of plasticine, is a useful technique to evaluate predation rates by vertebrates and invertebrates (Howe et al. 2009). Such techniques allow the predation rate of a given prey to be estimated by determining the number of predation marks as well as the identities of the predator groups based on distinctive bite marks. Distinctive marks can be attributed to chewing arthropods, small rodents, reptilians or birds (Barbaro et al. 2016).

In a more holistic way, it is often more informative to measure the potential biological control of a natural enemy community (parasitoids and predators) using different methods at the same time. To do this, researchers have used cage experiments and applied differential exclusion treatments (Rusch et al. 2013). Such approaches have the ability to evaluate the overall level of biological control in a given field over a given time period. Recently, a method using sentinel prey has been used to quantify the total biological control of natural enemies in vineyards (Rusch et al. 2017b). The exposure of grape moth eggs to parasitoids and predators is a rather easy technique that requires an insect stock culture with sufficient egg production. Fresh grape moth eggs can thus be exposed either on waxed paper or plastic sheets, but a more sophisticated procedure is to force females to oviposit on grape cuttings (either foliar or fructiferous) and to install such plants within the grape rows.

Table 3 Main bat species observed in Burgundy vineyards with recorded sonic casting flight or buzz. Observations and records by Gilles Sentenac (unpublished data)

Species	English common name	Casting flight	Buzz
<i>Pipistrellus pipistrellus</i>	Common pipistrelle bat	x	x
<i>Pipistrellus kuhlii</i>	Kuhl's pipistrelle bat	x	x
<i>Eptesicus serotinus</i>	Serotine bat	x	x
<i>P. kuhlii/nathusii</i>	Kuhl's/Nathusius' pipistrelle bat	x	x
<i>Nyctalus leisleri</i>	Leisler's bat	x	x
<i>Myotis myotis/blytii</i>	Greater/lesser mouse-eared bat	x	
<i>Nyctalus noctula</i>	Common noctule bat	x	x
<i>Plecotus</i> sp.	Brown/gray long-eared bat	x	
<i>Barbastella barbastellus</i>	Western barbastelle bat	x	x
<i>Myotis nattereri</i>	Natterer's bat	x	
<i>Myotis</i> sp.		x	
<i>Myotis mystacinus</i>	Whiskered bat	x	x
<i>Miniopterus schreibersii</i>	Schreibers' bent-winged bat	x	
<i>Myotis emarginatus</i>	Geoffroy's bat	x	
<i>Myotis bechsteinii</i>	Bechstein's bat	x	

Finally, some interesting in situ field video recordings were attempted in Australia to determine how habitat strata, and thus natural enemy communities, control another important polyphagous grape moth pest in Australian viticulture, *E. postvittana* (Frank et al. 2007). Similar techniques could be developed to measure the strength and efficiency of each member of such communities, and to evaluate the connectivity between the natural enemy habitat strata of predators and the grape clusters to be protected.

5 Key factors that induce variation in the biological control of grapevine moths

5.1 Effect of the host plant (including grape cultivar)

Phytophagous insect pests have evolved within a multitrophic environment, and many aspects of their physiology, behavior, and ecology have been shaped by interactions with other trophic levels (Poppy 1997; Karimzadeh and Wright 2008) (Fig. 3). Mainly, host plants directly affect the growth and adult fertility rates of phytophagous insects and indirectly the level of successful parasitism (Karimzadeh and Wright 2008). For example, a nutrient deficiency or/and toxic defensive compounds of the host plant slow-down the development of phytophagous insects, which extends the window of attacks for natural enemies (Benrey and Denno 1997). In grape moths, the egg size, the hatching success, and the larvae's developmental time depend on the grape variety. For instance, grape moths feeding on Gewürztraminer grapes have larger eggs, higher hatching success, and longer developmental times compared with moths feeding on other grape varieties (Moreau et al. 2006a, b). Moreover, the grape variety also

influences the egg and larval parasitism rates (Moreau et al. 2009, 2010). Overall, the egg parasitism is higher on cultivars on which *L. botrana* laid larger eggs, but some grape varieties, such as Pinot, appear to be somehow repellent to parasitoids (Moreau et al. 2009). Looking deeper into the trophic interactions in a vineyard is a promising way to improve the efficiency of biocontrol programs because it aids in understanding biological control failures.

The efficiency of the natural enemies depends on their ability to bypass the sophisticated defenses used by grapevine moths (Greeney et al. 2012). The first line of defense that is particularly efficient against larval parasitoids involves escaping, twisting, and dropping (Greeney et al. 2012; Vogelweith et al. 2014). Once this barrier is passed by the natural enemies, the cuticles represent a very efficient way to fight against these natural enemies. Indeed, an individual with a thick and resistance cuticle will be less likely to be parasitized by a parasitoid (Vogelweith et al. 2014). Once morphological and behavioral defenses are bypassed, the last and most efficient defense against natural enemies, such as parasitoids, is the immune system (Greeney et al. 2012) (Fig. 4). In insects, the immune system relies on constitutive and inducible mechanisms (Lavigne and Strand 2002; Siva-jothy et al. 2005). The constitutive defenses mostly involve the coordinated actions of the immune cells (the hemocytes) and the enzyme phenoloxidase (PO). Hemocytes insure most immune processes, such as the recognition and encapsulation of parasitoids and pathogens (Lavigne and Strand 2002; Cerenius and Söderhäll 2004; Siva-jothy et al. 2005), while PO mediates the melanization of foreign bodies (Fig. 3). Then, the induced response mainly consists of the production of specific antimicrobial peptides by the hemocytes and the fat bodies a few hours after an infection (Haine et al. 2008) (Fig. 3). This production of




Fig. 4 Summary diagram of the immune response of an insect larva after the injection of a parasitoid egg (F. Vogelweith). Once the parasitoid egg is injected, hemocytes will immediately recognize the foreign body via the recognition patterns at the surfaces. The immune reaction will start with the recruitment of other hemocytes to cover the egg. Hemocytes will

also allow the activation of the PO-PPO cascade which will release melanin to cover the egg and produce cytotoxic compounds to kill it. Few hours after the infection, antimicrobial peptides will be produced by the hemocyte to kill the remaining pathogens in the hemolymph

example, *E. ambiguella* invests more in physical defenses (thick and resistant integument) and less in behavioral defenses (e.g., twisting and dropping) and respond quickly to an immune challenge relative to *L. botrana* (Vogelweith et al. 2014). In the field, both the parasitism rate and successful parasitism are lower in *E. ambiguella* compared with *L. botrana* (Vogelweith et al. 2014), indicating that parasitoids might be more efficient in controlling *L. botrana* than *E. ambiguella* populations.

In the field, the larval immune system of *L. botrana* also depends on parasitism pressure (Vogelweith et al. 2013). Indeed, in populations experiencing a high parasitism pressure, larvae invest in higher immune parameter levels compared with populations under low parasitism pressure (Vogelweith et al. 2013). In populations with high levels of immune defenses, biological control might be difficult because their immune defenses could be adapted to the local high pressure from natural enemies.

These studies clearly show the value of increasing and applying our knowledge of tritrophic interactions between the host plant, the pests, and their natural enemies to improve the biological control of these insect pests. However, biological control was thought to be similar for *L. botrana* and *E. ambiguella* when co-occurring in vineyards. Based on their different defense-related investments, it could be important to integrate these results into biological control programs to select the most efficient natural enemies for each grapevine moth species.

5.2 Viticulture practices and biological control

Habitat manipulation is important for enhancing the biological control of arthropod pests. It is now well demonstrated that increasing vegetation diversity and/or structural complexity

within fields or in the close vicinity can decrease pest attacks, increase the influence of natural enemies, and reduce crop damage (Letourneau et al. 2011; Tonhasca and Byrne 1994). These effects can be explained by the natural enemy and the resource concentration hypotheses (Root 1973). The former suggests that natural enemies are more abundant and/or diverse, as well as more effective, in suppressing herbivore populations in more diverse plant communities (Andow 1991). This effect is attributed to the higher attractiveness of diverse habitats to predators as a result of an increased availability of resources. The latter suggests that reduced herbivore populations and plant damage levels in more diverse habitats occur because of the lower probability of herbivores to find their host plants (Otway et al. 2005; Plath et al. 2012). This effect is attributed to the chemical stimuli masking host plant odors, visual camouflage, and/or physical barriers limiting the movements of individuals (Finch and Collier 2000 and the references therein). For instance, releasing “odor-masking” substances by nonhost plant species confers some protection to the associated host plant in a diversified plant community (Thiéry and Visser 1986; Finch and Collier 2000).

Several studies have found that such processes operated in grapevine agroecosystems. First, cover crop management affects natural enemy communities and biological pest control. Increasing the within-field plant diversity enhances natural enemy abundance and/or diversity (Letourneau et al. 2009; Shields et al. 2016). Using indigenous grass cover crops within vineyards benefits natural enemies, such as generalist predators and parasitoids, and increases the biological control of tortricids (Danne et al. 2010). In a recent study, Rusch et al. (2017b) found that full grass cover within vineyards strongly limited the attack rates of tortricids compared with vineyards with partial grass cover, despite the effect of landscape context

on biological control. Altieri and Nicholls (2002) found higher infestation rates of two grape herbivores, including *L. botrana* in monocultures, when comparing different cropping systems (traditional vineyards based on agroforestry and modernized simple monoculture). They also found that monoculture vineyards exhibited lower numbers of predator and parasitoid species. Moreover, the provision of floral resources within vineyards by manipulating the ground cover increases the longevity and fecundity of parasitoids, which may result in higher parasitism rates of tortricids (Berndt et al. 2002; Begum et al. 2006; Berndt et al. 2006). For instance, buckwheat (*Fagopyrum esculentum*) and alyssum (*Lobularia maritima*) have beneficial effects on tortricid parasitoids (Begum et al. 2006). Similarly, mulching (e.g., covering the soil surface with a layer of organic material) enhances the abundance and/or diversity of carabids, hymenopteran parasitoids, dipteran parasitoids, hemipterans, and spiders without increasing insect pest abundance in vineyards (Thomson and Hoffmann 2007; Bruggisser et al. 2010). Second, contrary to such beneficial practices, tillage (e.g., reducing habitat complexity) has a negative effect on beneficial arthropods in vineyards (Sharley et al. 2008) as in other systems (Thorbeck and Bilde 2004). In addition, pesticide applications also explain the variability in the efficiency of tortricid biological control by impacting natural enemy communities. For instance, Nash et al. (2010) found an effect of season-long pesticide applications on arthropod community structure and showed that high pesticide metric scores negatively impact the activity levels of beneficial taxa, reducing the potential for biological control.

Beyond specific farming practices, organic systems promote the abundance and diversity levels of weeds and natural enemies (Bengtsson et al. 2005; Garratt et al. 2011; Tuck et al. 2014). Studies comparing organic and conventional farming in viticulture showed contrasting effects on natural enemies and biological pest control. For instance, Bruggisser et al. (2010) showed that organic farming promoted neither diversity nor abundance at any trophic level (plants, grasshoppers, and spiders) in Swiss vineyards. However, in intensive agricultural landscapes, organic farming favors a local plant species' richness (Nascimbene et al. 2012), and higher abundance and diversity levels of spiders and carabids have been found in organic farming compared with conventional farming (Gaigher and Samways 2010, 2014; Caprio et al. 2015). Despite a limited number of studies and some contradictory results, 6 out of 11 studies found on the Institute for Scientific Information's Web of Knowledge (currently Clarivate Analytics' Web of Science) reported that organic farming, compared with conventional farming, in vineyards tended to enhance the abundance and diversity levels of natural enemies (Caprio et al. 2015; Franin et al. 2016; Froidevaux et al. 2017; Geiger et al. 2010; Peverieri et al. 2009; Puig-Montserrat et al. 2017).

However, further research is needed to understand the context-dependent effects of organic farming. For instance,

relatively few studies have simultaneously examined the effects of organic and conventional farming on pest communities and crop damage, and nothing is known about the effective level of pest control between organic and conventional farming. Thus, even if pest species benefit from organic farming, they would exert a higher pressure on grape production even if the level of natural pest control is higher. In Bordeaux vineyards, for instance, grape moths were less parasitized in organic farming systems than in conventional ones (Rusch et al. 2015). This apparently counterintuitive result indicates that more studies examine the local and landscape effects of organic farming on natural enemies, biological control, and pest damage.

5.3 Vineyard landscapes and natural biological control

Landscape context, both in terms of composition and configuration, as well as trophic interactions, in agricultural systems affects population and community dynamics (Tscharntke et al. 2007; Rusch et al. 2010). Recent meta-analyses highlighted that landscape simplification through the loss of seminatural habitats affects the abundance and diversity levels of natural enemies as well as the biological control level (Chaplin-Kramer et al. 2011; Rusch et al. 2016). The positive effect of seminatural habitats on natural enemies and biological pest control is that they provide key resources and functions for natural enemies, such as overwintering sites and refuges from disturbances, alternative hosts or prey, as well as sources of nectar or pollen (Rusch et al. 2010). For instance, several natural enemies overwinter in woody or grassy habitats (Sarhou et al. 2014). Therefore, the spatial distribution between overwintering habitats and crops determines the distribution of individuals in the landscape (Rand et al. 2006). These results strongly suggest that conservation biological control will benefit from a landscape perspective.

Several studies found that maintaining seminatural habitats within vineyard-dominated landscapes benefits natural enemies and the biological control of tortricid moths (Thomson and Hoffmann 2009, 2013; Thomson et al. 2010; Barbaro et al. 2017; Pithon et al. 2016). The mean abundance of natural enemies within a vineyard declines with the distance from woody vegetation, leading to the higher parasitism and predation levels of tortricid moths in vine rows closer to woody habitats (Thomson and Hoffmann 2013). This distance effect was detected up to 40 m from the woody vegetation, thus providing guidelines for landscape management to increase pest control services. In a recent study, Barbaro et al. (2017) found that habitat heterogeneity at both local and landscape scales influences avian insectivory in vineyards by interacting with the avian community structure. Foliage-gleaning insectivores were found to be more abundant in landscapes that supported more seminatural habitats, suggesting an increase

in their contribution to pest control along the gradient of landscape complexity.

In conclusion, local and landscape complexity levels, as well as their interactions, are key drivers of natural enemy communities and the biological control of tortricid moths in vineyard-dominated landscapes. However, we still lack a good mechanistic understanding of the relationships between the structure of natural enemy communities (e.g., taxonomic or functional) and the level of biological control. Thus, further studies combining data and molecular analyses are needed to identify the roles of trophic assemblages and to highlight the optimal structure of the food web that promotes the biological pest control of tortricid pests.

5.4 Effects of climate changes on tritrophic interactions and natural biological control

As ectotherms, insect pests are greatly sensitive to environmental temperature, which regulates metabolic reactions (Kingsolver 2009) and influences many physiological processes, such as growth (Angilletta et al. 2004), development (Zuo et al. 2012), reproduction (Fischer et al. 2003), and immune functions (Murdock et al. 2012). Such a temperature dependency is tightly linked to the ecology and evolution of life histories (Ragland and Kingsolver 2008) and, therefore, must be considered when attempting to develop sustainable biocontrol strategies against grapevine moths, especially in a changing environment. Global warming and the expected 0.3–4.8 °C increase in mean global surface temperature at the end of this century underscore the importance of this idea (Intergovernmental Panel on Climate Change 2014). This rapidly changing climatic context can be linked to the “hotter-is-better” hypothesis to predict how GBMs will impact vineyards under warmer environmental conditions. Indeed, this hypothesis postulates that ectothermic species adapted to warm conditions display the highest maximal fitness level and reach this optimum at high environmental temperatures (Kingsolver 2009; Knies et al. 2009). Thus, global warming benefits GBM by bringing the environmental temperature closer to their thermal optimum, especially in areas where thermal conditions remain colder than their thermal optimum (for instance, northern areas of the distribution range) (Deutsch et al. 2008). Thus, global warming may result in high GBM growth rates (Knies et al. 2009) and greater damage levels to crops. In addition to a direct thermal effect on GBM, global warming might also indirectly affect their performance by influencing two associated trophic levels: grapevine (Jones and Davis 2000) and natural enemies, like parasitoids (see Hance et al. 2007; Reineke and Thiéry 2016 for reviews). We summarize here several impacts of climate change on the physiology of grapevine moths and its consequences in terms of yield losses and pest management strategies.

Like many insects, *L. botrana* exhibits a higher developmental rate when exposed to warmer conditions (Torres-Villa 1996; Martín-Vertedor et al. 2010). Additionally, temperature is an environmental factor that triggers the termination of diapause in this species (Roditakis and Karandinos 2001). As a consequence, mild early springs promote a significant advance in the first emergence wave of adults from overwintering pupae and impact the voltinism of this species (Martín-Vertedor et al. 2010; Reineke and Thiéry 2016). For instance, in 2006, warmer conditions allowed the emergence of a fourth generation of *L. botrana* in Mediterranean areas, even though this species is usually trivoltine (Martín-Vertedor et al. 2010). This additional generation likely intensified the pressure exerted on crops (see Caffarra et al. 2012 and references therein). However, grapes also respond to global warming by shifting to an earlier phenology (e.g., earlier bud burst and blooming), which results in a significant advance in harvest dates and a shortened crop life cycle. This is observed in France where winegrowers are currently collecting grape bunches almost 1 month earlier than during the last 50 years (Jones and Davis 2000; Seguin and de Cortazar 2005; Schultz and Jones 2010). By doing so, they remove the resources that potentially allow the complete development of the late-season generation of the pests and their overwintering as diapausing pupae (Martín-Vertedor et al. 2010; Caffarra et al. 2012; Reineke and Thiéry 2016). Despite these potential benefits of global warming for the control of GBM populations, harvest dates are not just determined by temperature but also by grapevine variety, which influences plant phenology (Jones and Davis 2000). Consequently, it could be hypothesized that early ripening varieties, like Chardonnay, would be less prone to the detrimental effects of an additional GBM generation than late ripening varieties that would be exposed to the late pest generation before harvest (Caffarra et al. 2012). The detrimental effects of an additional generation on vine production could, therefore, be modulated by grape cultivars and the corresponding plant phenology.

Faster GBM development may influence their interactions with enemies naturally present or released in vineyards (see Section 3) and the associated consequences on biological control strategies that rely on these natural enemies. Temperature is a decisive factor modulating the ability of pests to fight against parasitoids through behavioral or their immune responses. To our knowledge, no study to date has investigated the direct effects of contrasting thermal conditions on the ability of GBMs to resist parasitoid pressure. However, recent results suggest that GBMs' immune functions can be modulated in the field by temperature (Vogelweith et al. 2013). In the scenario of a rising global temperature, the increased immune functions of grapevine pests would be problematic because they enhance the pests' ability to resist natural enemies, especially parasitoids, which decreases the effectiveness of biocontrol programs.

6 Perspectives and conclusions

The body of knowledge we have reviewed highlights several key factors operating at multiple spatial scales that influence the biological control of grapevine moths. We particularly show that host plant, farming practices at the field scale, and landscape context, as well as climate change, are driving grapevine moth population dynamics through direct effects on pest populations or indirect effects mediated by their natural enemies. This review also provides a framework for understanding the context dependency of the efficiency of biological control programs targeted against these insect pests.

6.1 Challenges for research

Several research avenues addressing the biological control of grapevine moths clearly need further investigation. This includes the diversity of macroorganisms that participate in the biological control of grape pests. In addition to parasitoids, spiders and harvestmen are clearly important components of predator webs. However, their biology, behavior, and more generally, life history traits should be further investigated. Progress in barcoding techniques should also allow deeper investigations into the functional ecology of food webs and into who eats whom and at what intensity level. Our understanding of these parameters is increasing in several agrosystems, including vineyards, in which there is a particular focus on predators of grape berry moths. This research should provide rapid progress that is accessible to grape growers. Understanding factors that influence host/prey quality is also crucial for avoiding unsuspected failures in biological control. Research on the roles played by the pest host plant should therefore be intensified for the induction of pest resistance, including the immune system expression, to biocontrol agents. Additionally, in our opinion, the possible attraction and arrestment roles of maternal pheromones (see Section 3) should receive more attention. The regulation of natural enemies by landscape factors, including plant biodiversity and especially floral interrow plantings, is now being studied in vineyards. These efforts should be intensified, especially identifying which types of flowers provide the best fitness gains for parasitoids and predators.

6.2 Challenges for growers and crop protection advisors

The main new finding that can be quickly implemented to help protect grapes from grape moths is the installation of grass or flower strips in vineyards to favor natural enemy activity. The reduction of insecticide use is now in progress in most European grape-producing countries, and this will favor the use of biological controls. Control practices, like the use of mating disruption or *B. thuringiensis*, should also be developed because limited harmful effects on biodiversity

have been reported. Very recent results also indicate that the organic production of grape favors increased biodiversity and the capability of natural control (Muneret et al. 2017). These practices could be quickly implemented in viticulture.

All of the studies we reviewed illustrated that many factors influence the efficiency of biological control. Additionally, the impact of global warming on *L. botrana* is complex and thus hard to predict, being dependent on coevolutionary plant–pest and pest–parasitoid mechanisms in the context of tritrophic interactions. Deeper insights into these coevolutionary mechanisms in the face of climate change will be needed to improve our predictions of the future impact of pests and, subsequently, the use of biocontrol strategies. In addition, we emphasized the need for improved knowledge of trophic relationships between natural enemies and GBM in vineyards to predict the level of tortricid control. Determining the roles of the different species involved in grape moth regulation, as well as considering the whole community, is a major challenge in enhancing biological control in actual and future vineyard systems. The reduction of insecticide consumption in European viticulture requires that obstacles to biocontrol be addressed and that several biological control strategies be developed.

Acknowledgements We thank Dr. Lesley Benyon, PhD, from Edanz Group for editing a final draft of this manuscript and two anonymous reviewers for improving the manuscript.

Funding information The two PhD students (LM and CI) were respectively funded by a French Fund for Research on Biodiversity (FRB) Région Nouvelle Aquitaine and a French Ministry of Research RT grants. The French Casdar Project Biocontrôle as well as the SOLUTION project (FRB) funded some results presented here. We also thank the Conseil Régional of Bourgogne (FABER).

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

References

- Abrol DP, Shankar U (2012) History, overview and principles of ecologically based pest management. Integrated pest management: principles and practice. ABI Publishing, Wallingford. <https://doi.org/10.1079/9781845938086.0001>
- Agusti N, Bourguet D, Spataro T, Delos M, Eychenne N, Folcher L, Arditi R (2005) Detection, identification and geographical distribution of European corn borer larval parasitoids using molecular markers. *Mol Ecol* 14:3267–3274. <https://doi.org/10.1111/j.1365-294X.2005.02650.x>
- Altieri MA, Nicholls CI (2002) The simplification of traditional vineyard based agroforests in northwestern Portugal: some ecological implications. *Agrofor Syst* 56:185–191
- Andow DA (1991) Vegetational diversity and arthropod population response. *Annu Rev Entomol* 36:561–586

- Angilletta MJ, Steury TD, Sears MW (2004) Temperature, growth rate, and body size in ectotherms: fitting pieces of a life-history puzzle. *Integr Comp Biol* 44:498–509. <https://doi.org/10.1093/icb/44.6.498>
- Audouin V (1842) Histoire des insectes nuisibles de la vigne et particulièrement de la pyrale. Fortin Masson et cie, Paris
- Bael SAV, Philpott SM, Greenberg R, Bichier P, Barber NA, Mooney KA, Gruner DS (2008) Birds as predators in tropical agroforestry systems. *Ecology* 89:928–934. <https://doi.org/10.1890/06-1976.1>
- Barbaro L, Rusch A, Muirir E, Gravelier B, Thiery D, Castagnyrol B (2017) Avian pest control in vineyards is driven by interactions between bird functional diversity and landscape heterogeneity. *J Appl Ecol* 54:500–508. <https://doi.org/10.1111/1365-2664.12740>
- Barnay O, Pizzol J, Gertz C, Kienlen JC, Hommay G, Lapchin L (1999) Host density-dependence of discovery and exploitation rates of egg patches of *Lobesia botrana* (Lepidoptera: Tortricidae) and *Ephesia kunhiella* (Lepidoptera: Pyralidae) by the parasitoid *Trichogramma cacoeciae* (Hymenoptera: Trichogrammatidae). *J Econ Entomol* 92: 1311–1320
- Begum M, Gurr GM, Wratten SD, Hedberg PR, Nicol HI (2006) Using selective food plants to maximize biological control of vineyard pests. *J Appl Ecol* 43:547–554. <https://doi.org/10.1111/j.1365-2664.2006.01168.x>
- Bengtsson J, Ahnström J, Weibull AC (2005) The effects of organic agriculture on biodiversity and abundance: a meta-analysis. *J Appl Ecol* 42:261–269. <https://doi.org/10.1111/j.1365-2664.2005.01005.x>
- Benrey B, Denno RF (1997) The slow-growth-high mortality hypothesis: a test using the cabbage butterfly. *Ecology* 78:987–999
- Berndt LA, Wratten SD, Hassan PG (2002) Effects of buckwheat flowers on leafroller (Lepidoptera: Tortricidae) parasitoids in a New Zealand vineyard. *Agric For Entomol* 4:39–45
- Berndt LA, Wratten SD, Scarratt SL (2006) The influence of floral resource subsidies on parasitism rates of leafrollers (Lepidoptera: Tortricidae) in New Zealand vineyards. *Biol Control* 37:50–55. <https://doi.org/10.1016/j.biocontrol.2005.12.005>
- Birkhofer K, Bylund H, Dalin P, Ferlian O, Gagic V, Hambäck PA, Klapwijk M, Mestre L, Roubinet E, Schroeder M, Stenberg JA, Porcel M, Bjökman C, Jonsson M (2017) Methods to identify the prey of invertebrate predators in terrestrial field studies. *Ecol Evol* 7: 1942–1953. <https://doi.org/10.1002/ece3.2791>
- Bolduc E, Buddle CM, Bostanian NJ, Vincent C (2005) Ground-dwelling spider fauna (Araneae) of two vineyards in southern Quebec. *Environ Entomol* 34:635–645. <https://doi.org/10.1603/0046-225X-34.3.635>
- Boreau de Roince C, Lavigne C, Ricard JM, Franck P, Bouvier JC, Garcin A, Symondson WO (2012) Predation by generalist predators on the codling moth versus a closely-related emerging pest the oriental fruit moth: a molecular analysis. *Agric For Entomol* 14:260–269. <https://doi.org/10.1111/j.1461-9563.2011.00564.x>
- Boyer S, Zhang H, Lempérière G (2012) A review of control methods and resistance mechanisms in stored-product insects. *Bull Entomol Res* 102:213–229. <https://doi.org/10.1017/S0007485311000654>
- Brewer MJ, Goodell PB (2012) Approaches and incentives to implement integrated pest management that addresses regional and environmental issues. *Annu Rev Entomol* 57:41–59. <https://doi.org/10.1146/annurev-ento-120709-144748>
- Bruggisser OT, Schmidt-Entling MH, Bacher S (2010) Effects of vineyard management on biodiversity at three trophic levels. *Biol Conserv* 143:1521–1528. <https://doi.org/10.1016/j.biocon.2010.03.034>
- Butault J-P, Dedryver C-A, Gary C, Guichard L, Jacquet F, Meynard JM, Nicot P, Pitrat M, Reau R, Sauphanor B, Savini I, Volay T (2010) Synthèse du rapport d'étude Écophyto R&D: quelles voies pour réduire l'usage des pesticides? INRA Editions. Hal: hal-01172967
- Caffarra A, Rinaldi M, Eccel E, Rossi V, Pertot I (2012) Modelling the impact of climate change on the interaction between grapevine and its pests and pathogens: European grapevine moth and powdery mildew. *Agric Ecosyst Environ* 148:89–101. <https://doi.org/10.1016/j.agee.2011.11.017>
- Caprio E, Nervo B, Isaia M, Allegro G, Rolando A (2015) Organic versus conventional systems in viticulture: comparative effects on spiders and carabids in vineyards and adjacent forests. *Agric Syst* 136:61–69. <https://doi.org/10.1016/j.agsy.2015.02.009>
- Cerenius L, Söderhäll K (2004) The prophenoloxidase-activating system in invertebrates. *Immunol Rev* 198:116–126. <https://doi.org/10.1111/j.0105-2896.2004.00116.x>
- Chaplin-Kramer R, O'Rourke ME, Blitzer EJ, Kremen C (2011) A meta-analysis of crop pest and natural enemy response to landscape complexity. *Ecol Lett* 14:922–932. <https://doi.org/10.1111/j.1461-0248.2011.01642.x>
- Charbonnier Y, Barbaro L, Theillout A, Jactel H (2014) Numerical and functional responses of forest bats to a major insect pest in pine plantations. *PLoS One* 9(10):e109488. <https://doi.org/10.1371/journal.pone.0109488>
- Chuche J, Xuéreb A, Thiéry D (2006) Attraction of *Dibrachys cavus* (Hymenoptera: Pteromalidae) to its host frass volatiles. *J Chem Ecol* 32:2721–2731. <https://doi.org/10.1007/s10886-006-9195-8>
- Cozzi G, Pascale M, Perrone G, Visconti A, Logrieco A (2006) Effect of *Lobesia botrana* damages on black *aspergilli*'s rot and ochratoxin A content in grapes. *Int J Food Microbiol* 111:88–92. <https://doi.org/10.1016/j.ijfoodmicro.2006.03.012>
- Crowder DW, Northfield TD, Strand MR, Snyder WE (2010) Organic agriculture promotes evenness and natural pest control. *Nature* 466: 109–112. <https://doi.org/10.1038/nature09183>
- Danne A, Thomson LJ, Sharley DJ, Penfold CM, Hoffmann AA (2010) Effects of native grass cover crops on beneficial and pest invertebrates in Australian vineyards. *Environ Entomol* 39:970–978. <https://doi.org/10.1603/EN09144>
- Delbac L, Thiéry D (2016) Damage to grape flowers and berries by *Lobesia botrana* larvae (Denis & Schiffemüller) (Lepidoptera: Tortricidae) and relation to larval age. *Aust J Grape Wine Res* 22(2):256–261. <https://doi.org/10.1111/ajgw.12204>
- Desneux N, Decourtye A, Delpuech JM (2007) The sublethal effects of pesticides on beneficial arthropods. *Annu Rev Entomol* 52:81–106. <https://doi.org/10.1146/annurev.ento.52.110405.091440>
- Dhaliwal GS, Jindal V, Dhawan AK (2010) Insect pest problems and crop losses: changing trends. *Indian J Ecol* 37:1–7
- Ehler LE (2006) Integrated pest management (IPM): definition, historical development and implementation, and the other IPM. *Pest Manag Sci* 62:787–789
- Endure (2006). Pesticide use in viticulture, available data on current practices and innovations, bottlenecks and need for research in this field and specific leaflets analysing the conditions of adoption of some innovations, EC project deliverable DR1.23, 35p
- Fatouros NE, Dicke M, Mumm R, Meiners T, Hilker M (2008) Foraging behaviour of egg parasitoids exploiting chemical information. *Behav Ecol* 19:677–689. <https://doi.org/10.1093/beheco/arn011>
- Ferguson HM, John B, Ng'habi K, Knols BG (2005) Redressing the sex imbalance in knowledge of vector biology. *Trends Ecol Evol* 20: 202–209. <https://doi.org/10.1016/j.tree.2005.02.003>
- Fischer K, Brakefield PM, Zwaan BJ (2003) Plasticity in butterfly egg size: why larger offspring at lower temperatures? *Ecology* 84:3138–3147. <https://doi.org/10.1890/02-0733>
- Franin K, Barić B, Kuštera G (2016) The role of ecological infrastructure on beneficial arthropods in vineyards. *Span J Agric Res* 14:303–311. <https://doi.org/10.5424/sjar/2016141-7371>
- Frank SD, Wratten SD, Sandhu HS, Shrewsbury PM (2007) Video analysis to determine how habitat strata affects predator diversity and predation of *Epiphyas postvittana* (Lepidoptera: Tortricidae) in a vineyard. *Biol Control* 41(2):230–236
- Frenoy C, Durier C, Hawlitzky N (1992) Effect of kairomones from egg and female-adult stages of *Ostrinia nubilalis* Hubner (Lepidoptera:

- Pyalidae) on *Trichogramma brassicae* Bedzdenko (Hymenoptera: Trichogrammatidae) female kinesis. *J Chem Ecol* 18(5):761–773
- Froidevaux JSP, Louboutin B, Jones G (2017) Does organic farming enhance biodiversity in Mediterranean vineyards? A case study with bats and arachnids. *Agric Ecosyst Environ* 249:112. <https://doi.org/10.1016/j.agee.2017.08.012>
- Gabel B, Thiéry D (1992) Biological evidence of an oviposition deterring pheromone in *Lobesia botrana* (Lepidoptera: Tortricidae). *J Chem Ecol* 18:353–358
- Gabel B, Thiéry D (1994) Non-host plant odour (*Tanacetum vulgare*; Asteraceae) affects the reproductive behaviour of *Lobesia botrana* Den. et Schiff (Lepidoptera: Tortricidae). *J Insect Behav* 7:149–157
- Gabel B, Thiéry D (1996) Oviposition response of *Lobesia botrana* females to long-chain free fatty acids and esters from its eggs. *J Chem Ecol* 22:161–171
- Gaigher R, Samways MJ (2010) Surface-active arthropods in organic vineyards, integrated vineyards and natural habitat in the Cape Floristic Region. *J Insect Conserv* 14:595–605. <https://doi.org/10.1007/s10841-010-9286-2>
- Gaigher R, Samways MJ (2014) Landscape mosaic attributes for maintaining ground-living spider diversity in a biodiversity hotspot. *Insect Conserv Div* 7:470–479. <https://doi.org/10.1111/icad.12070>
- Garratt MPD, Wright DJ, Leather SR (2011) The effects of farming system and fertilisers on pests and natural enemies: a synthesis of current research. *Agric Ecosyst Environ* 141:261–270. <https://doi.org/10.1016/j.agee.2011.03.014>
- Geiger F, Bengtsson J, Berendse F, Weisser WW, Emmerson M, Morales MB, Onate JJ, Eggers S, Pärt T, Bommarco R, Bengtsson L, Clement LW, Weisser WW, Olszewski A, Ceryngier P, Hawro V, Inchausti P, Fischer C, Flohre A, Thies C, Tscharntke T (2010) Persistent negative effects of pesticides on biodiversity and biological control potential on European farmland. *Basic Appl Ecol* 11:97–105. <https://doi.org/10.1016/j.baae.2009.12.001>
- Genini M (2000) Faune épigée de la vigne et des milieux environnants. *Rev Suisse Vitic Arboric Hortic* 32:1–12
- Glen DM (1977) Predation of codling moth eggs, *Cydia pomonella*, the predators responsible and their alternative prey. *J Appl Ecol* 14(2): 445–456
- Glen DM, Milsom NF (1978) Survival of mature larvae of codling moth (*Cydia pomonella*) on apple trees and ground. *Ann Appl Biol* 90: 133–146
- Glenn DC, Hoffmann AA (1997) Developing a commercially viable system for biological control of light brown apple moth (Lepidoptera: Tortricidae) in grapes using endemic *Trichogramma* (Hymenoptera: Trichogrammatidae). *J Econ Entomol* 90:370–338
- Goulet H, Lesage L, Bostanian NJ, Vincent C, Lasnier J (2004) Diversity and seasonal activity of ground beetles (Coleoptera: Carabidae) in two vineyards of southern Quebec, Canada. *Ann Entomol Soc Am* 97(6):1263–1272
- Greeney HF, Dyer L, Smilanich M (2012) Feeding by lepidopteran larvae is dangerous: a review of caterpillars' chemical, physiological, morphological, and behavioral defenses against natural enemies. *Invertebr Surviv J* 9:7–34
- Haine ER, Moret Y, Siva-Jothy MT, Rolff J (2008) Antimicrobial defense and persistent infection in insects. *Science* 322:1257–1259. <https://doi.org/10.1126/science.1165265>
- Hallenbeck WH, Cunningham-Burns KM (1995) Pesticides and human health. University of Illinois, Springer, Chicago
- Hance T, van Baaren J, Vernon P, Boivin G (2007) Impact of extreme temperatures on parasitoids in a climate change perspective. *Annu Rev Entomol* 52:107–126. <https://doi.org/10.1146/annurev.ento.52.110405.091333>
- Hogg BN, Daane KM (2010) The role of dispersal from natural habitat in determining spider abundance and diversity in California vineyards. *Agric Ecosyst Environ* 135:260–267. <https://doi.org/10.1016/j.agee.2009.10.004>
- Hommay G, Gertz C, Kienlen JC, Pizzol J, Chavigny P (2002) Comparison between the control efficacy of *Trichogramma evanescens* Westwood (Hymenoptera: Trichogrammatidae) and two *Trichogramma cacoeciae*. *Biocontrol Sci Tech* 12(5):569–581
- Hommay G, Kienlen JC, Gertz C, Bihri C, Pizzol J (2011) Factors affecting the post release dispersal of *Trichogramma cacoeciae* Marchal in the vineyards. *IOBC/WPRS Bull* 67:207–214
- Howe A, Lövei GL, Nachman G (2009) Dummy caterpillars as a simple method to assess predation rates on invertebrates in a tropical agroecosystem. *Entomol Exp Appl* 131:325–329. <https://doi.org/10.1111/j.1570-7458.2009.00860.x>
- Hrcek JAN, Miller SE, Quicke DL, Smith M (2011) Molecular detection of trophic links in a complex insect host–parasitoid food web. *Mol Ecol Res* 11:786–794. <https://doi.org/10.1111/j.1755-0998.2011.03016.x>
- Intergovernmental Panel on Climate Change (2014) Climate change 2014: impacts, adaptation, and vulnerability. Part A: global and sectoral aspects. Contributions of working group II to the fifth assessment report of the intergovernmental panel on climate change. Cambridge University Press, Cambridge. <https://doi.org/10.1017/CBO9781107415379>
- Ioriatti C, Anfora G, Tasin M, de Cristofaro A, Witzgall P, Lucchi A (2011) Chemical ecology and management of *Lobesia botrana* (Lepidoptera: Tortricidae). *J Eco Entomol* 104:1125–1137. <https://doi.org/10.1603/EC10443>
- Jedlicka JA, Greenberg R, Letourneau DK (2011) Avian conservation practices strengthen ecosystem services in California vineyards. *PLoS One* 6:e27347. <https://doi.org/10.1371/journal.pone.0027347>
- Jedlicka JA, Letourneau DK, Cornelisse TM (2014) Establishing songbird nest boxes increased avian insectivores and reduced herbivorous arthropods in a Californian vineyard, USA. *Conserv Evid* 11: 34–38
- Jolicoeur H (1894) Les ravageurs de la vigne. Michaud publication, Reims
- Jones GV, Davis RE (2000) Climate influences on grapevine phenology, grape composition, and wine production and quality for Bordeaux, France. *Am J Viti Enol* 51:249–261
- Jourdie V, Alvarez N, Turlings TCJ (2008) Identification of seven species of hymenopteran parasitoids of *Spodoptera frugiperda*, using polymerase chain reaction amplification and restriction enzyme digestion. *Agric For Entomol* 10(2):129–136. <https://doi.org/10.1111/j.1461-9563.2008.00362.x>
- Karimzadeh J, Wright DJ (2008) Bottom-up cascading effects in a tritrophic system: interactions between plant quality and host-parasitoid immune responses. *Ecol Entomol* 33:45–52. <https://doi.org/10.1111/j.1365-2311.2007.00933.x>
- Karp DS, Mendenhall CD, Sandi RF, Chaumont N, Ehrlich PR, Hadly EA, Daily GC (2013) Forest bolsters bird abundance, pest control and coffee yield. *Ecol Lett* 16:1339–1347. <https://doi.org/10.1111/ele.12173>
- Kingsolver JG (2009) ASN Presidential Address: the well-temperated biologist. *Am Nat* 174:755–768
- Knies JL, Kingsolver JG, Burch CL (2009) Hotter is better and broader: thermal sensitivity of fitness in a population of bacteriophages. *Am Nat* 173:419–430. <https://doi.org/10.1086/597224>
- Kogan M (1998) Integrated pest management: historical perspectives and contemporary developments. *Annu Rev Entomol* 43:243–270
- Lavine MD, Strand MR (2002) Insect hemocytes and their role in immunity. *Insect Biochem Mol Biol* 32:1295–1309
- Lazzaro BP, Little TJ (2009) Immunity in a variable world. *Philos Trans R Soc Lond Ser B Biol Sci* 364:15–26
- Letourneau DK, Jedlicka JA, Bothwell SG, Moreno CR (2009) Effects of natural enemy biodiversity on the suppression of arthropod herbivores in terrestrial ecosystems. *Annu Rev Ecol Evol Syst* 40:573–592. <https://doi.org/10.1146/annurev.ecolsys.110308.120320>

- Letourneau DK, Armbrrecht I, Rivera BS, Lerma JM, Carmona EJ, Daza MC, Escobar S, Galindo V, Gutiérrez C, López SD, Mejía JL, Rangel AMA, Rangel JH, Rivera L, Saavedra CA, Torres AM, Trujillo AR (2011) Does plant diversity benefit agroecosystems? A synthetic review. *Ecol Appl* 21:9–21. <https://doi.org/10.1890/09-2026.1>
- Loni A, Samartsev KG, Scaramozzino PL, Belokobylskij SA, Lucchi A (2016) Braconinae parasitoids (Hymenoptera, Braconidae) emerged from larvae of *Lobesia botrana* (Denis & Schiffmüller) (Lepidoptera, Tortricidae) feeding on *Daphne gnidium* L. *ZooKeys* 587:125–150. <https://doi.org/10.3897/zookeys.587.8478>
- Loreau M, Hector A (2001) Partitioning selection and complementarity in biodiversity experiments. *Nature* 412:72–76. <https://doi.org/10.1038/35083573>
- Maas B, Clough Y, Tschamtké T (2013) Bats and birds increase crop yield in tropical agroforestry landscapes. *Ecol Lett* 16:1480–1487. <https://doi.org/10.1111/ele.12194>
- Macfadyen S, Gibson R, Polaszek A, Morris RJ, Craze PG, Planque R, Symondson WO, Memmott J (2009) Do differences in food web structure between organic and conventional farms affect the ecosystem service of pest control? *Ecol Lett* 12:229–238. <https://doi.org/10.1111/j.1461-0248.2008.01279.x>
- Marchal P, Feytaud J (1911) Sur un parasite des œufs de la cochylys et de l'eudémis. *C R Acad Sci* 153:633–635
- Martin EA, Reineking B, Seo B, Steffan-Dewenter I (2013) Natural enemy interactions constrain pest control in complex agricultural landscapes. *Proc Natl Acad Sci U S A* 110:5534–5539. <https://doi.org/10.1073/pnas.1215725110>
- Martín-Vertedor D, Ferrero-García J, Torres-Vila LM (2010) Global warming affects phenology and voltinism of *Lobesia botrana* in Spain. *Agric For Entomol* 12:169–176. <https://doi.org/10.1111/j.1461-9563.2009.00465.x>
- Mills NJ, Danne KM (2005) Biological and cultural controls. Non pesticide alternative can suppress crop pests. *Calif Agric* 59:23–28
- Milonas PG, Martinou AF, Kontamida DC, Karamaouna F, Kostantopoulou (2009) Attraction of different trichogramma species to parys olea sex pheromone. *Ann Entomol Soc Amer* 102:1145–1150
- Moreau J, Benrey B, Thiéry D (2006a) Assessing larval food quality for phytophagous insects: are the facts as simple as they appear? *Funct Ecol* 20:592–600. <https://doi.org/10.1111/j.1365-2435.2006.01145.x>
- Moreau J, Benrey B, Thiéry D (2006b) Grape variety affects larval performance and also female reproductive performance of the European grapevine moth *Lobesia botrana* (Lepidoptera: Tortricidae). *Bull Entomol Res* 96:205–212. <https://doi.org/10.1079/BER2005417>
- Moreau J, Richard A, Benrey B, Thiéry D (2009) Host plant cultivar of the grapevine moth *Lobesia botrana* affects the life history traits of an egg parasitoid. *Biol Control* 50:117–122. <https://doi.org/10.1016/j.biocontrol.2009.03.017>
- Moreau J, Villemant C, Benrey B, Thiéry D (2010) Species diversity and composition of larval parasitoids of the European grapevine moth (*Lobesia botrana*): the influence of region and cultivar. *Biol Control* 54:300–306. <https://doi.org/10.1016/j.biocontrol.2010.05.019>
- Moreau J, Monceau K, Thiéry D (2016) Larval food influences oviposition dynamics and egg quality traits in females of *Lobesia botrana*. *J Pest Sci* 89:439–448
- Moret Y, Schmid-Hempel P (2000) Survival for immunity: the price of immune system activation for bumblebee workers. *Science* 290:1166–1168
- Muller K, Vogelweith F, Thiéry D, Moret Y, Moreau J (2015) Immune benefits from alternative host plants could maintain polyphagy in a phytophagous insect. *Oecologia* 177:467–475. <https://doi.org/10.1007/s00442-014-3097-1>
- Muneret L, Thiéry D, Joubard B, Rusch A (2017) Deployment of organic farming at landscape scale maintains low pest infestation and high crop productivity in vineyards. *J Appl Ecol* 1–10. doi: <https://doi.org/10.1111/1365-2664.13034>
- Murdoch CC, Paaijmans KP, Cox-Foster D, Read AF, Thomas MB (2012) Rethinking vector immunology: the role of environmental temperature in shaping resistance. *Nat Rev Microbiol* 10:869–876. <https://doi.org/10.1038/nrmicro2900>
- Nagarkatti S, Tobin C, Saunders MC, Muza AJ (2003) Release of native *Trichogramma minutum* to control grapeberry moth. *Can Entomol* 135:589–598. <https://doi.org/10.4039/n02-099>
- Naranjo SE, Ellsworth PC (2009) Fifty years of the integrated control concept: moving the model and implementation forward in Arizona. *Pest Manag Sci* 65:1267–1286. <https://doi.org/10.1002/ps.1861>
- Nascimbene J, Marini L, Paoletti MG (2012) Organic farming benefits local plant diversity in vineyard farms located in intensive agricultural landscapes. *Environ Manag* 49:1054–1060. <https://doi.org/10.1007/s00267-012-9834-5>
- Nash MA, Hoffmann AA, Thomson LJ (2010) Identifying signature of chemical applications on indigenous and invasive nontarget arthropod communities in vineyards. *Ecol Appl* 20:1693–1703
- Oerke EC (2006) Crop losses to pests. *J Agri Sci* 144:31–43
- Otway SJ, Hector A, Lawton JH (2005) Resource dilution effects on specialist insect herbivores in a grassland biodiversity experiment. *J Anim Ecol* 74:234–240
- Papura D, Rusch A, Roux P, Delbac L, Thiéry D (2016) Early detection and identification of larval parasitoids in *Lobesia botrana* using PCR-RFLP method. *Biol Control* 103:95–100
- Paull C, Austin AD (2007) The hymenopteran parasitoids of light brown apple moth, *Epiphyas postvittana* (Walker) (Lepidoptera: Tortricidae) in Australia. *Aust J Entomol* 45(2006):142–156. <https://doi.org/10.1111/j.1440-6055>
- Pérez-Staples D, Shelly TE, Yuval B (2012) Female mating failure and the failure of ‘mating’ in sterile insect programs. *Entomol Exp Appl* 146:66–78
- Pertot I, Caffi T, Mugnai L, Hoffmann C, Grando MS, Gary C, Lafond D, Duso C, Thiéry D, Mazzoni V, Anfora G (2017) A critical review of plant protection tools for reducing pesticide use on grapevine and new perspectives for implementation of IPM in viticulture. *Crop Protec* 97:70–84. <https://doi.org/10.1016/j.cropro.2016.11.025>
- Peverieri GS, Simoni S, Goggioli D, Liguori M, Castagnoli M (2009) Effects of variety and management practices on mite species diversity in Italian vineyards. *Bull Insectol* 62(1):53–60
- Pithon JA, Beaujouan V, Daniel H, Pain G, Vallet J (2016) Are vineyards important habitats for birds at local or landscape scales? *Basic Appl Ecol* 17:240–251
- Plath M, Dom S, Riedel J, Barrios H, Mody K (2012) Associational resistance and associational susceptibility: specialist herbivores show contrasting responses to tree stand diversification. *Oecologia* 169:477–487
- Ponton F, Wilson K, Cotter SC, Raubenheimer D, Simpson SJ (2011) Nutritional immunology: a multi-dimensional approach. *PLoS Pathog* 7:e1002223
- Poppy G (1997) Tritrophic interactions: improving ecological understanding and biological control? *Endeavour* 21:61–65
- Puig-Montserrat X, Stefanescu C, Torre I, Palet J, Fàbregas E, Dantart J, Arrizabalaga A, Flaquer C (2017) Effects of organic and conventional crop management on vineyard biodiversity. *Agric Ecosyst Environ* 243:19–26
- Ragland GJ, Kingsolver JG (2008) The effect of fluctuating temperatures on ectotherm life-history traits: comparisons among geographic populations of *Wyeomyia smithii*. *Evol Ecol Res* 10:29–44
- Rand TA, Tylisanakis JM, Tschamtké T (2006) Spillover edge effects: the dispersal of agriculturally subsidized insect natural enemies into adjacent natural habitats. *Ecol Lett* 9:603–614. <https://doi.org/10.1111/j.1461-0248.2006.00911.x>

- Reda Abd el Monsef AI (2004) Biological control of grape berry moths *Eupoecilia ambiguella* HB and *Lobesia botrana* Den. et Schiff. (Lepidoptera Tortricidae) by using egg parasitoids of the genus *Trichogramma*. Dissertation, Universität Giesen
- Reineke A, Thiéry D (2016) Grapevine insect pests and their natural enemies in the age of global warming. *J Pest Sci* 89:313–328. <https://doi.org/10.1007/s10340-016-0761-8>
- Roberts MJ (2009) Guide des araignées de France et d'Europe: Plus de 450 Espèces Décrites et Illustrées. Delachaux et Niestlé
- Roditakis NE, Karandinos MG (2001) Effects of photoperiod and temperature on pupal diapause induction of grape berry moth *Lobesia botrana*. *Physiol Entomol* 26:329–340
- Root RB (1973) Organization of a plant-arthropod association in simple and diverse habitats: the fauna of collards (*Brassica oleracea*). *Ecol Monogr* 43:95–124
- Roush R, Tabashnik BE (1990) Pesticide resistance in arthropods. Chapman and Hall, London
- Rusch A, Valantin-Morison M, Sarthou JP, Roger-Estrade J (2010) Biological control of insect pests in agroecosystems: effects of crop management, farming systems, and seminatural habitats at the landscape scale: a review. *Adv Agron* 109:219–259. [https://doi.org/10.1016/S0065-2113\(10\)09006-1](https://doi.org/10.1016/S0065-2113(10)09006-1)
- Rusch A, Bommarco R, Jonsson M, Smith HG, Ekbom B (2013) Flow and stability of natural pest control services depend on complexity and crop rotation at the landscape scale. *J Appl Ecol* 50:345–354. <https://doi.org/10.1111/1365-2664.12055>
- Rusch A, Delbac L, Muneret L, Thiéry D (2015) Organic farming and host density affect parasitism rates of tortricid moths in vineyards. *Agr Ecosyst Environ* 214:46–53. <https://doi.org/10.1016/j.agee.2015.08.019>
- Rusch A, Chaplin-Kramer R, Gardiner MM, Hawro V, Holland J, Landis D, Thies C, Tschantke T, Weisser WW, Winqvist C (2016) Agricultural landscape simplification reduces natural pest control: a quantitative synthesis. *Agr Ecosyst Environ* 221:198–204. <https://doi.org/10.1016/j.agee.2016.01.039>
- Rusch A, Bommarco R, Ekbom B (2017a) Conservation biological control in agricultural landscapes. In: Sauvion N, Calatayud PA, Thiéry D (eds) Insect plant interactions in a crop protection perspective. *Adv Bot Res* 81:333–360. <https://doi.org/10.1016/bs.abr.2016.11.001>
- Rusch A, Delbac L, Thiéry D (2017b) Grape moth density in Bordeaux vineyards depends on local habitat management despite effects of landscape heterogeneity on their biological control. *J Appl Ecol* 54:1794–1803. <https://doi.org/10.1111/1365-2664.12858>
- Rydell J, Entwistle A, Racey PA (1996) Timing and foraging flights of three species of bats in relation to insect activity and predation risk. *Oikos* 76:243–252
- Sarthou JP, Badoz A, Vaissière B, Chevallier A, Rusch A (2014) Local more than landscape parameters structure natural enemy communities during their overwintering in semi-natural habitats. *Agric Ecosyst Environ* 194:17–28. <https://doi.org/10.1016/j.agee.2014.04.018>
- Schellhorn NA, Gagic V, Bommarco R (2015) Time will tell: resource continuity bolsters ecosystem services. *Trends Ecol Evol* 30:524–530. <https://doi.org/10.1016/j.tree.2015.06.007>
- Schultz HR, Jones GV (2010) Climate induced historic and future changes in viticulture. *J Wine Res* 21:137–145. <https://doi.org/10.1080/09571264.2010.530098>
- Seguin B, de Cortazar IG (2005) Climate warming: consequences for viticulture and the notion of ‘terroirs’ in Europe. *Acta Hort* 689:61–71. <https://doi.org/10.17660/ActaHortic.2005.689.3>
- Sentenac G (2011) La faune auxiliaire des vignobles de France. Editions France Agricole, Paris
- Sharley DJ, Hoffmann AA, Thomson LJ (2008) The effects of soil tillage on beneficial invertebrates within the vineyard. *Agric For Entomol* 10:233–243. <https://doi.org/10.1111/j.1461-9563.2008.00376.x>
- Shields MW, Tompkins J, Saville DJ, Meurk CD, Wratten S (2016) Potential ecosystem service delivery by endemic plants in New Zealand vineyards: successes and prospects. *PeerJ* 4:e2042. <https://doi.org/10.7717/peerj.2042>
- Siva-Jothy MT, Moret Y, Rolff J (2005) Insect immunity: an evolutionary ecology perspective. *Adv Insect Physiol* 32:1–48. [https://doi.org/10.1016/S0065-2806\(05\)32001-7](https://doi.org/10.1016/S0065-2806(05)32001-7)
- Skopura JP, Hothem RL (1985) Consumption of commercially-grown grapes by american robins: a field evaluation of laboratory estimates. *J Field Ornithol* 56(4):369–378
- Speakman JR, Rydell J (2000) Avoidance behaviour of bats and moths: when is it predator defence? *Oikos* 88:221–223
- Straub CS, Finke DL, Snyder WE (2008) Are the conservation of natural enemy biodiversity and biological control compatible goals? *Biol Control* 45:225–237
- Thacker JRM (2002) An introduction to arthropod pest control. Cambridge University Press, Cambridge
- Thiéry D (2008) Les tordeuses nuisibles à la vigne. In: Esmenjaud D, Kreiter S, Martinez M, Sforza R, Thiéry D, van Helden M, Yvon M (eds) Les ravageurs de la vigne. Féret Publication, pp 214–246
- Thiéry D (2011) Gaps in knowledge for modern integrated protection in viticulture: lessons from controlling grape berry moths. *IOBC/WPRS Bull* 67:305–311
- Thiéry D, Visser JH (1986) Masking of host plant odour in the olfactory orientation of the Colorado potato beetle. *Entomol Exp Appl* 41:165–172
- Thiéry D, Gabel B, Farkas P, Jarry M (1995) Egg dispersion in codling moth: influence of egg extract and of its fatty acid constituents. *J Chem Ecol* 21:2015–2026
- Thiéry D, Xuéreb A, Villemant C, Sentenac G, Delbac L, Kuntzman P (2001) Les parasites de tordeuses de vignobles : aperçu de quelques espèces présentes dans 3 régions viticoles françaises. *IOBC/WPRS Bull* 24(7):135–141
- Thiéry D, Yoshida T, Guisnet M (2006) *Phytomyptera nigrina* (Meigen) (Diptera, Tachinidae) parasite of the first generation of the European grapevine moth larvae in several vineyards of the Roussillon area. *Tachinid Times* 19:1–4
- Thiéry D, Carton Y, Vidal C, Gauthier N, Derridj S, Vercambre B, Goebel F-R, Grégoire J-C, Lieutier F (2011a) Histoire de l'installation de quelques ravageurs. In: Sauvion N, Calatayud PA, Thiéry D, Marion-Poll F (eds) Interactions Insectes plantes. Quae-ird Publication, pp 623–673
- Thiéry D, Delbac L, Villemant C, Moreau J (2011b) Control of grape berry moth larvae using parasitoids: should it be developed? *IOBC/WPRS Bull* 67:189–196
- Thiéry D, Monceau K, Moreau J (2014) Larval intraspecific competition for food in the European grapevine moth *Lobesia botrana*. *Bull Entomol Res* 104:517–524. <https://doi.org/10.1017/S0007485314000273>
- Thomas MB (1999) Ecological approaches and the development of “truly integrated” pest management. *Proc Nat Acad Sci* 96:5944–5951
- Thomson LJ, Hoffmann AA (2007) Effects of ground cover (straw and compost) on the abundance of natural enemies and soil macro invertebrates in vineyards. *Agric For Entomol* 9:173–179. <https://doi.org/10.1111/j.1461-9563.2007.00322.x>
- Thomson LJ, Hoffmann AA (2009) Vegetation increases the abundance of natural enemies in vineyards. *Biol Control* 49:259–269. <https://doi.org/10.1016/j.biocontrol.2009.01.009>
- Thomson LJ, Hoffmann AA (2013) Spatial scale of benefits from adjacent woody vegetation on natural enemies within vineyards. *Biol Control* 64:57–65. <https://doi.org/10.1016/j.biocontrol.2012.09.019>
- Thomson LJ, Macfadyen S, Hoffmann AA (2010) Predicting the effects of climate change on natural enemies of agricultural pests. *Biol Control* 52:296–306. <https://doi.org/10.1016/j.biocontrol.2009.01.022>

- Thorbeck P, Bilde T (2004) Reduced numbers of generalist arthropod predators after crop management. *J Appl Ecol* 41:526–538. <https://doi.org/10.1111/j.0021-8901.2004.00913.x>
- Tilman D, Balzer C, Hill J, Befort BL (2011) Global food demand and the sustainable intensification of agriculture. *Proc Nat Acad Sci* 108:20260–20264. <https://doi.org/10.1073/pnas.1116437108>
- Tonhasca A, Byrne DN (1994) The effects of crop diversification on herbivorous insects: a meta-analysis approach. *Ecol Entomol* 19:239–244
- Torres-Vila LM, Stockel J, Roehrich R, Rodriguez-Molina MC (1997) The relation between dispersal and survival of *Lobesia botrana* larvae and their density in vine inflorescences. *Entomol Exp Appl* 84:109–114
- Torres-Villa LM (1996) Efecto de la temperatura de desarrollo preimaginal sobre el potencial biótico de la pollila del racimo de la vid, *Lobesia botrana* (Denis y Schiffermüller, [1775]) (Lepidoptera: Tortricidae). *SHILAP Revta Lepid* 24:197–206
- Traugott M, Zangerl P, Juen A, Schallhart N, Pfiffner L (2006) Detecting key parasitoids of lepidopteran pests by multiplex PCR. *Biol Control* 39:39–46. <https://doi.org/10.1016/j.biocontrol.2006.03.001>
- Tscharntke T, Bommarco R, Clough Y, Crist TO, Kleijn D, Rand TA, Tylianakis JM, van Nouhuys S, Vidal S (2007) Conservation biological control and enemy diversity on a landscape scale. *Biol Control* 43:294–309. <https://doi.org/10.1016/j.biocontrol.2007.08.006>
- Tuck SL, Winqvist C, Mota F, Ahnström J, Turnbull LA, Bengtsson J (2014) Land-use intensity and the effects of organic farming on biodiversity: a hierarchical meta-analysis. *J Appl Ecol* 51:746–755. <https://doi.org/10.1111/1365-2664.12219>
- Unruh TR, Miliczky ER, Horton DR, Thomsen-Archer K, Rehfield-Ray L, Jones VP (2016) Gut content analysis of arthropod predators of codling moth in Washington apple orchards. *Biol Control* 102:85–92. <https://doi.org/10.1016/j.biocontrol.2016.05.014>
- Vincent C, Isaacs R, Bostanian NJ, Lasnier J (2012) Principles of arthropod pest management in vineyards. In: Bostanian NJ, Vincent C, Isaacs R (eds) *Arthropod management in vineyards: pests, approaches and future directions*. Springer, Dordrecht, pp 1–15. https://doi.org/10.1007/978-94-007-4032-7_1
- Vogelweith F, Thiéry D, Quaglietti B, Moret Y, Moreau J (2011) Host plant variation plastically impacts different traits of the immune system of a phytophagous insect. *Funct Ecol* 25:1241–1247. <https://doi.org/10.1111/j.1365-2435.2011.01911>
- Vogelweith F, Dourneau M, Thiéry D, Moret Y, Moreau J (2013) Geographical variation in parasitism shapes larval immune function in a phytophagous insect. *Naturwissenschaften* 100:1149–1161. <https://doi.org/10.1007/s00114-013-1119-1>
- Vogelweith F, Thiéry D, Moret Y, Colin E, Motreuil S, Moreau J (2014) Defense strategies used by two sympatric vineyard moth pests. *J Insect Physiol* 64:54–61. <https://doi.org/10.1016/j.jinsphys.2014.03.009>
- Vogelweith F, Moreau J, Thiéry D, Moret Y (2015) Food-mediated modulation of immunity in a phytophagous insect: an effect of nutrition rather than parasitic contamination. *J Insect Physiol* 77:55–61. <https://doi.org/10.1016/j.jinsphys.2015.04.003>
- Vogelweith F, Moret Y, Monceau K, Thiéry D, Moreau J (2016) The relative abundance of hemocyte types in a polyphagous moth larva depends on diet. *J Insect Physiol* 88:33–39. <https://doi.org/10.1016/j.jinsphys.2016.02.010>
- Walter GH (2005) *Insect pest management and ecological research*. Cambridge University Press, Cambridge
- Walton VM, Daane K, Addison P (2012) Biological control of arthropods and its application in vineyards. In: Bostanian NJ, Vincent C, Isaacs R (eds) *Arthropod management in vineyards: pests, approaches and future directions*. Springer, Dordrecht, pp 91–117. https://doi.org/10.1007/978-94-007-4032-7_5
- Way MJ, Van Emden HF (2000) Integrated pest management in practice—pathways towards successful application. *Crop Prot* 19:81–103
- Wirta HK, Hebert PDN, Kaartinen R, Prosser SW, Várkonyi G, Roslin T (2014) Complementary molecular information changes our perception of food web structure. *Proc Nat Acad Sci* 111:1885–1890
- Xuéreb A, Thiéry D (2006) Does natural larval parasitism of *Lobesia botrana* vary between years, generation, density of the host and vine cultivar? *Bull Entomol Res* 96:105–110. <https://doi.org/10.1079/BER2005393>
- Zuo W, Moses ME, West GB, Hou C, Brown JH (2012) A general model for effects of temperature on ectotherm ontogenetic growth and development. *Proc R Soc B* 79:1840–1846. <https://doi.org/10.1098/rspb.2011.2000>